

Addenda/Corrigenda

Fixed Points and Two-cycles of the Discrete Logarithm

Joshua Holden

Department of Mathematics, Rose-Hulman Institute of Technology, Terre Haute, IN,
47803-3999, USA, holden@rose-hulman.edu

Abstract. This report consists of additions and corrections to the author's paper [1], which appeared in the proceedings of the ANTS V conference. The work described here was presented at the conference itself, which took place after the original paper was published. The abstract of the original paper was as follows: We explore some questions related to one of Brizolis: does every prime p have a pair (g, h) such that h is a fixed point for the discrete logarithm with base g ? We extend this question to ask about not only fixed points but also two-cycles. Campbell and Pomerance have not only answered the fixed point question for sufficiently large p but have also rigorously estimated the number of such pairs given certain conditions on g and h . We attempt to give heuristics for similar estimates given other conditions on g and h and also in the case of two-cycles. These heuristics are well-supported by the data we have collected, and seem suitable for conversion into rigorous estimates in the future.

The formulas leading up to Conjecture 7 have several typos. The corrected formulas should read:

$$\sum_{m|p-1} |S_m|^2 / |T_m| \approx \sum_{m|p-1} \frac{\phi(m)}{m^2} \left(\sum_{d|(p-1)/m} \frac{\phi(dm)}{d} \right)^2$$

$$N_{(3),h \text{ ANY}, a \text{ ANY}}(p) \approx (p-1) + \sum_{m|p-1} \frac{\phi(m)}{m^2} \left(\sum_{d|(p-1)/m} \frac{\phi(dm)}{d} \right)^2$$

(This is also the formula which should appear in Conjecture 7(a).)

$$\sum_{m|p-1} \frac{\phi(m)^3}{m^2} \left(\sum_{d|(p-1)/m} \frac{\phi(d)}{d} \right)^2 = \sum_{m|p-1} \left(\prod_{q|m} \frac{\phi(q)^3}{q^2} \right) \left(\prod_{q|(p-1)/m} \left(1 + \frac{\phi(q)}{q} \right)^2 \right)$$

$$= \prod_{q|p-1} \left(\frac{\phi(q)^3}{q^2} + \left(1 + \frac{\phi(q)}{q} \right)^2 \right) = \prod_{q|p-1} \left(q + 1 - \frac{1}{q} \right)$$

In addition, there are two errors in (4); the equation should read:

$$\begin{aligned} \sum_{m|p-1} |S_m|^2 / |T_m| &\approx \prod_q \left(\sum_{\beta=0}^{\alpha} \phi(q^\beta) \left[\left(1 - \frac{1}{q} \right) (\alpha - \beta) + \frac{\phi(q^\beta)}{q^\beta} \right]^2 \right) \\ &= \prod_q \left(\left[\left(1 - \frac{1}{q} \right) \alpha + 1 \right]^2 + \sum_{\beta=1}^{\alpha} q^\beta \left(1 - \frac{1}{q} \right) \left[\left(1 - \frac{1}{q} \right) (\alpha - \beta + 1) \right]^2 \right) \\ &= \prod_q \left(\left[\left(1 - \frac{1}{q} \right) \alpha + 1 \right]^2 \right. \\ &\quad \left. + \left(1 - \frac{1}{q} \right)^3 \left[(\alpha + 1)^2 \frac{q^{\alpha+1} - q}{q - 1} - 2(\alpha + 1) \frac{\alpha q^{\alpha+2} - (\alpha + 1)q^{\alpha+1} + q}{(q - 1)^2} \right. \right. \\ &\quad \left. \left. + \frac{\alpha^2 q^{\alpha+3} - (2\alpha^2 + 2\alpha - 1)q^{\alpha+2} + (\alpha^2 + 2\alpha + 1)q^{\alpha+1} - q^2 - q}{(q - 1)^3} \right] \right) \quad (4) \end{aligned}$$

Conjecture 1(c) is incorrect. In (1) if h PR then g PR also, so $N_{(1),g \text{ ANY},h \text{ PR}}(p)$ is in fact equal to $N_{(1),g \text{ PR},h \text{ RPPR}}(p)$.

The same observation for (2) gives $N_{(2),g \text{ PR},h \text{ RPPR}}(p) = N_{(2),g \text{ ANY},h \text{ RPPR}}(p)$ in Conjecture 5 and $N_{(2),g \text{ PR},h \text{ PR}}(p) = N_{(2),g \text{ ANY},h \text{ PR}}(p)$ in Conjecture 6(b).

In Conjectures 2 and 6(a) it should be noted that the observed values in question must be exactly equal, by the symmetry of (3). Likewise in Conjectures 4 and 6(b).

A complete and corrected list of Theorems and Conjectures follows. The numbering from the original paper has been preserved as much as possible.

Proposition 1. $N_{(1),g \text{ ANY},h \text{ RP}}(p) = \phi(p - 1)$.

Theorem 1 (Zhang, independently by others).

$$\begin{aligned} N_{(1),g \text{ PR},h \text{ RPPR}}(p) &= N_{(1),g \text{ PR},h \text{ RP}}(p) \\ &= N_{(1),g \text{ PR},h \text{ PR}}(p) \\ &= N_{(1),g \text{ ANY},h \text{ RPPR}}(p) \\ &= N_{(1),g \text{ ANY},h \text{ PR}}(p) \\ &\approx \phi(p - 1)^2 / (p - 1) \end{aligned}$$

Conjecture 1.

- (a) $N_{(1),g \text{ ANY},h \text{ ANY}}(p) \approx p - 1$
- (b) $N_{(1),g \text{ PR},h \text{ ANY}}(p) \approx \phi(p - 1)$
- (c) (See above.)
- (d) $N_{(1),g \text{ RP},h \bullet}(p) \approx \phi(p - 1) / (p - 1) N_{(1),g \text{ ANY},h \bullet}(p)$

$$(e) \ N_{(1),g \text{RPPR},h\bullet}(p) \approx \phi(p-1)/(p-1)N_{(1),g \text{PR},h\bullet}(p)$$

Conjecture 2.

$$\begin{aligned} N_{(2),g \text{ANY},h \text{RP}}(p) &= N_{(3),h \text{RP},a \text{ANY}}(p) \\ &\approx 2\phi(p-1). \end{aligned}$$

Conjecture 3.

$$\begin{aligned} N_{(2),h \text{RP},g \text{ORD } h}(p) &= N_{(3),h \text{RP},a \text{RP}}(p) \\ &\approx \phi(p-1) + \phi(p-1)^2/(p-1). \end{aligned}$$

Conjecture 4.

$$\begin{aligned} N_{(2),g \text{PR},h \text{RP}}(p) &= N_{(3),h \text{RP},a \text{PR}}(p) \\ &\approx 2\phi(p-1)^2/(p-1). \end{aligned}$$

Conjecture 5.

$$\begin{aligned} N_{(2),g \text{PR},h \text{RPPR}}(p) &= N_{(2),g \text{ANY},h \text{RPPR}}(p) \\ &= N_{(3),h \text{RPPR},a\bullet}(p) \\ &= N_{(3),h\bullet,a \text{RPPR}}(p) \\ &= N_{(3),h \text{RPPR},a \text{RPPR}}(p) \\ &\approx \phi(p-1)^2/(p-1) + \phi(p-1)^3/(p-1)^2. \end{aligned}$$

Conjecture 6.

(a)

$$\begin{aligned} N_{(2),h \text{ANY},g \text{ORD } h}(p) &= N_{(3),h \text{ANY},a \text{RP}}(p) \\ &\approx 2\phi(p-1). \end{aligned}$$

(b)

$$\begin{aligned} N_{(2),g \text{PR},h \text{PR}}(p) &= N_{(2),g \text{ANY},h \text{PR}}(p) \\ &= N_{(3),h \text{PR},a \text{RP}}(p) \\ &\approx 2\phi(p-1)^2/(p-1). \end{aligned}$$

Conjecture 7.

- (a) $N_{(3),h \text{ANY},a \text{ANY}}(p) \approx (p-1) + \sum_{m|p-1} \frac{\phi(m)}{m^2} \left(\sum_{d|(p-1)/m} \frac{\phi(dm)}{d} \right)^2$.
- (b) If $p-1$ is squarefree then $N_{(3),h \text{ANY},a \text{ANY}}(p) \approx (p-1) + \prod_{q|p-1} \left(q + 1 - \frac{1}{q} \right)$, where the product is taken over primes q dividing $p-1$.
- (c) In general, $N_{(3),h \text{ANY},a \text{ANY}}(p) \approx (p-1)$ plus the formula given in (4).
- (d) $N_{(3),h \text{PR},a \text{ANY}}(p) \approx 2\phi(p-1)$.
- (e) $N_{(3),h \text{ANY},a \text{PR}}(p) \approx 2\phi(p-1)$.

$$(f) \ N_{(3),h \text{ PR},a \text{ PR}}(p) \approx \phi(p-1) + \phi(p-1)^2/(p-1).$$

Conjecture 8.

$$(a) \ N_{(2),g \text{ PR},h \text{ ANY}}(p) \approx 2\phi(p-1).$$

$$(b) \ N_{(2),g \text{ ANY},h \text{ ANY}}(p) \approx 2(p-1).$$

Conjecture 9.

$$(a) \ N_{(2),g \text{ RP},h \bullet}(p) \approx \phi(p-1)/(p-1)N_{(2),g \text{ ANY},h \bullet}(p).$$

$$(b) \ N_{(2),g \text{ RPPR},h \bullet}(p) \approx \phi(p-1)/(p-1)N_{(2),g \text{ PR},h \bullet}(p).$$

These conjectures are summarized in Tables A-1, A-2, and A-3, which also contain new data presented at ANTS V. This data was collected on a Beowulf cluster, with 19 nodes, each consisting of 2 Pentium III processors running at 1 Ghz. The programming was done in C, using MPI, OpenMP, and OpenSSL libraries. The collection took 68 hours for all values of $N_{(\bullet),\bullet,\bullet}(p)$, for five primes p starting at 100000. Table A-4 summarizes the relationships between solutions to (2) and solutions to (3).

Table A-1. Solutions to (1)

(a) Predicted formulas for $N_{(1)}(p)$

$g \setminus h$	ANY	PR	RP	RPPR
ANY	$\approx_{(p-1)}$	$\approx \frac{\phi(p-1)^2}{(p-1)}$	$= \phi(p-1)$	$\approx \frac{\phi(p-1)^2}{(p-1)}$
PR	$\approx \phi(p-1)$	$\approx \frac{\phi(p-1)^2}{(p-1)}$	$\approx \frac{\phi(p-1)^2}{(p-1)}$	$\approx \frac{\phi(p-1)^2}{(p-1)}$
RP	$\approx \phi(p-1)$	$\approx \frac{\phi(p-1)^3}{(p-1)^2}$	$\approx \frac{\phi(p-1)^2}{(p-1)}$	$\approx \frac{\phi(p-1)^3}{(p-1)^2}$
RPPR	$\approx \frac{\phi(p-1)^2}{(p-1)}$	$\approx \frac{\phi(p-1)^3}{(p-1)^2}$	$\approx \frac{\phi(p-1)^3}{(p-1)^2}$	$\approx \frac{\phi(p-1)^3}{(p-1)^2}$

(b) Predicted values for $N_{(1)}(100057)$

$g \setminus h$	ANY	PR	RP	RPPR
ANY	100056	9139.46	30240	9139.46
PR	30240	9139.46	9139.46	9139.46
RP	30240	2762.23	9139.46	2762.23
RPPR	9139.46	2762.23	2762.23	2762.23

(c) Observed values for $N_{(1)}(100057)$

$g \setminus h$	ANY	PR	RP	RPPR
ANY	98506	9192	30240	9192
PR	29630	9192	9192	9192
RP	29774	2784	9037	2784
RPPR	9085	2784	2784	2784

Finally, the analysis in the last paragraph of Section 2 is incomplete. If $\gcd(h, a, p-1) = 1$, then the correspondence between solutions of (2) and solutions of (3) is one-to-one. (E.g., if $h \text{ RP}$ or $a \text{ RP}$.) If $\gcd(h, a, p-1) > 1$, however,

Table A-2. Solutions to (3)

(a) Predicted formulas for the nontrivial part of $N_{(3)}(p)$				
$a \setminus h$	ANY	PR	RP	RPPR
ANY	$\approx \sum \frac{ S_m ^2}{ T_m }$	$\approx \phi(p-1)$	$\approx \phi(p-1)$	$\approx \frac{\phi(p-1)^3}{(p-1)^2}$
PR	$\approx \phi(p-1)$	$\approx \frac{\phi(p-1)^2}{(p-1)}$	$\approx \frac{\phi(p-1)^2}{(p-1)}$	$\approx \frac{\phi(p-1)^3}{(p-1)^2}$
RP	$\approx \phi(p-1)$	$\approx \frac{\phi(p-1)^2}{(p-1)}$	$\approx \frac{\phi(p-1)^2}{(p-1)}$	$\approx \frac{\phi(p-1)^3}{(p-1)^2}$
RPPR	$\approx \frac{\phi(p-1)^3}{(p-1)^2}$	$\approx \frac{\phi(p-1)^3}{(p-1)^2}$	$\approx \frac{\phi(p-1)^3}{(p-1)^2}$	$\approx \frac{\phi(p-1)^3}{(p-1)^2}$

(b) Predicted values for the nontrivial part of $N_{(3)}(100057)$				
$a \setminus h$	ANY	PR	RP	RPPR
ANY	190822.0	30240	30240	2762.225
PR	30240	9139.458	9139.458	2762.225
RP	30240	9139.458	9139.458	2762.225
RPPR	2762.225	2762.225	2762.225	2762.225

(c) Observed values for the nontrivial part of $N_{(3)}(100057)$				
$a \setminus h$	ANY	PR	RP	RPPR
ANY	190526	30226	30291	2820
PR	30226	9250	9231	2820
RP	30291	9231	9086	2820
RPPR	2820	2820	2820	2820

more than one solution to (2) may give the same solution to (3). This will be explored in more detail in a forthcoming paper.

References

1. Joshua Holden. Fixed points and two-cycles of the discrete logarithm. In C. Fieker and D.R. Kohel, editors, *Algorithmic Number Theory (ANTS 2002)*, number 2369 in LNCS, pages 405–415. Springer, 2002. <http://link.springer-ny.com/link/service/series/0558/bibs/2369/23690405.htm>.

Table A-3. Solutions to (2)

(a) Predicted formulas for the nontrivial part of $N_{(2)}(p)$

$g \setminus h$	ANY	PR	RP	RPPR
ANY	$\approx_{(p-1)}$	$\approx_{\phi(p-1)}$	$\approx \frac{\phi(p-1)^2}{(p-1)}$	$\approx \frac{\phi(p-1)^3}{(p-1)^2}$
PR	$\approx_{\phi(p-1)}$	$\approx \frac{\phi(p-1)^2}{(p-1)}$	$\approx \frac{\phi(p-1)^2}{(p-1)}$	$\approx \frac{\phi(p-1)^3}{(p-1)^2}$
RP	$\approx_{\phi(p-1)}$	$\approx \frac{\phi(p-1)^2}{(p-1)}$	$\approx \frac{\phi(p-1)^3}{(p-1)^2}$	$\approx \frac{\phi(p-1)^4}{(p-1)^3}$
RPPR	$\approx \frac{\phi(p-1)^2}{(p-1)}$	$\approx \frac{\phi(p-1)^3}{(p-1)^2}$	$\approx \frac{\phi(p-1)^3}{(p-1)^2}$	$\approx \frac{\phi(p-1)^4}{(p-1)^3}$

(b) Predicted values for the nontrivial part of $N_{(2)}(100057)$

$g \setminus h$	ANY	PR	RP	RPPR
ANY	100056	9139.5	30240	2762.2
PR	30240	9139.5	9139.5	2762.2
RP	30240	2762.2	9139.5	834.8
RPPR	9139.5	2762.2	2762.2	834.8

(c) Observed values for the nontrivial part of $N_{(2)}(100057)$

$g \setminus h$	ANY	PR	RP	RPPR
ANY	100860	9231	30291	2820
PR	30850	9231	9231	2820
RP	30368	2882	9240	916
RPPR	9376	2882	2882	916

Table A-4. Relationship between solutions to (2) and solutions to (3)

$a \setminus h$	ANY	PR	RP	RPPR
ANY			g ANY h RP	g PR h RPPR
PR			g PR h RP	g PR h RPPR
RP	h ANY g ORD h	g PR h PR	h RP g ORD h	g PR h RPPR
RPPR	g PR h RPPR	g PR h RPPR	g PR h RPPR	g PR h RPPR